



SKIV2L gene

Ski2 like RNA helicase

Normal Function

The *SKIV2L* gene provides instructions for making a protein whose function has not been confirmed. Based on its similarity to a protein in other organisms, researchers speculate that the *SKIV2L* protein acts as part of a group of proteins called the SKI complex. This complex is thought to be necessary for the function of another large protein complex known as the cytosolic exosome. Within cells, the cytosolic exosome helps to recognize and break down excess or abnormal messenger RNA (mRNA) molecules. mRNA is a chemical cousin of DNA that serves as the genetic blueprint for protein production. Studies suggest that the cytosolic exosome's role in getting rid of excess and abnormal mRNA is important for cell growth.

Health Conditions Related to Genetic Changes


trichohepatoenteric syndrome

At least nine mutations in the *SKIV2L* gene have been found to cause trichohepatoenteric syndrome, a rare condition that affects many parts of the body. Its major signs and symptoms include chronic diarrhea starting in infancy, hair abnormalities, distinctive facial features, and liver disease. Mutations in this gene likely eliminate the function of the *SKIV2L* protein. Researchers hypothesize that a loss of this protein's function impairs the activity of the SKI complex and the cytosolic exosome. However, it is unknown how these changes could lead to chronic diarrhea and the other features of trichohepatoenteric syndrome.

Chromosomal Location

Cytogenetic Location: 6p21.33, which is the short (p) arm of chromosome 6 at position 21.33

Molecular Location: base pairs 31,958,804 to 31,969,852 on chromosome 6 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- 170A
- DDX13
- helicase-like protein
- helicase SKI2W
- HLP
- SKI2
- SKI2 homolog, superkiller viralicidic activity 2-like
- SKI2W
- SKIV2
- SKIV2L1
- superkiller viralicidic activity 2-like (*S. cerevisiae*)
- THES2

Additional Information & Resources

Educational Resources

- Genomes (second edition, 2002): Degradation of mRNAs
<https://www.ncbi.nlm.nih.gov/books/NBK21132/#A7399>
- Marie Curie Bioscience Database: Structural Components and Architectures of RNA Exosomes
<https://www.ncbi.nlm.nih.gov/books/NBK45033/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28SKIV2L%5BTIAB%5D%29+OR+%28SKI2W%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D>

OMIM

- SUPERKILLER VIRALICIDIC ACTIVITY 2, S. CEREVISIAE, HOMOLOG-LIKE
<http://omim.org/entry/600478>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_SKIV2L.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=SKIV2L%5Bgene%5D>
- HGNC Gene Family: RNA helicases
<http://www.genenames.org/cgi-bin/genefamilies/set/1168>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=10898
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/6499>
- UniProt
<http://www.uniprot.org/uniprot/Q15477>

Sources for This Summary

- Fabre A, Charroux B, Martinez-Vinson C, Roquelaure B, Odul E, Sayar E, Smith H, Colomb V, Andre N, Hugot JP, Goulet O, Lacoste C, Sarles J, Royet J, Levy N, Badens C. SKIV2L mutations cause syndromic diarrhea, or trichohepatoenteric syndrome. *Am J Hum Genet.* 2012 Apr;90(4):689-92. doi: 10.1016/j.ajhg.2012.02.009. Epub 2012 Mar 22.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22444670>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3322239/>
- Fabre A, Martinez-Vinson C, Goulet O, Badens C. Syndromic diarrhea/Tricho-hepato-enteric syndrome. *Orphanet J Rare Dis.* 2013 Jan;8:5. doi: 10.1186/1750-1172-8-5. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23302111>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3560276/>
- van Dijk EL, Schilders G, Pruijn GJ. Human cell growth requires a functional cytoplasmic exosome, which is involved in various mRNA decay pathways. *RNA.* 2007 Jul;13(7):1027-35. Epub 2007 Jun 1.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17545563>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1894934/>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/SKIV2L>

Reviewed: March 2014

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications

U.S. National Library of Medicine

National Institutes of Health

Department of Health & Human Services